

CHAPTER 1: INTRODUCTION TO VOLUME TWO

INTRODUCTION

Coastal habitats provide ecological, cultural, and economic value. They act as critical habitat for thousands of species, including numerous threatened and endangered species, by providing shelter, spawning grounds, and food (Mitsch and Gosselink 2000). They often act as natural buffers, providing ecological, social, and economic benefits by filtering sediment and pollution from upland drainage thereby improving water quality, reducing the effects of floodwaters and storm surges, and preventing erosion. In addition to these ecosystem services, healthy coastal habitats provide many human values including opportunities for:

- Outdoor recreation and tourism
- Education
- Traditional use and subsistence lifestyles
- Healthy fishing communities, and
- Obtaining other marketable goods

Therefore, healthy functioning coastal habitats are not only important ecologically, they also support healthy coastal communities and, more generally, improve the quality of human lives. Despite these benefits, coastal habitats have been modified, degraded, and removed throughout the United States and its protectorates beginning with European colonization (Dahl 1990). Thus, many coastal habitats around the United States are in desperate need of restoration and subsequent monitoring of restoration projects.

WHAT IS RESTORATION MONITORING?

The science of restoration requires two basic tools: the ability to manipulate ecosystems to recreate a desired community and the ability to evaluate whether the manipulation has produced the desired change (Keddy 2000). The latter is often referred to as restoration monitoring.

For this manual, restoration monitoring is defined as follows:

“The systematic collection and analysis of data that provides information useful for measuring project performance at a variety of scales (locally, regionally, and nationally), determining when modification of efforts are necessary, and building long-term public support for habitat protection and restoration.”

Restoration monitoring contributes to the understanding of complex ecological systems (Meeker et al. 1996) and is essential in documenting restoration performance and adapting project and program approaches when needs arise. If results of monitoring restored coastal areas are disseminated, they can provide tools for planning management strategies and help improve future restoration practices and projects (Washington et al. 2000). Restoration monitoring can be used to determine whether project goals are being met and if mid-course corrections are necessary. It provides information on whether selected project goals are good measures for future projects and how to perform routine maintenance in restored areas (NOAA et al. 2002). Monitoring also provides the basis for a rigorous review of the pre-construction project planning and engineering.

Restoration monitoring is closely tied to and directly derived from restoration project goals. The monitoring plan (i.e., what is measured, how often, when, and where) should be developed with project goals in mind. If, for example, the goal of a restoration project is to increase the amount of fish utilizing a coastal marsh, then measurements should be selected that can quantify progress toward that goal. A variety of questions about sampling techniques

and protocols need to be answered before monitoring can begin. For the fish utilization example, these may include:

- Will active or passive capture techniques be used (e.g., beach seines vs. fyke nets)?
- Where and when will samples be taken?
- Who will conduct the sampling?
- What level of identification will be required?
- What structural characteristics such as water level fluctuation or water chemistry will also be monitored and how?
- Who is responsible for housing and analyzing the data?
- How will results of the monitoring be disseminated?

Each of these questions, as well as many others, will be answered with the goals of the restoration project in mind. These questions need to be addressed before any measurements are taken in the field. In addition, although restoration monitoring is typically thought of as a ‘post-restoration’ activity, practitioners will find it beneficial to collect some data before and during project implementation. Pre-implementation monitoring provides baseline information to compare with post-implementation data to see if the restoration is having the desired effect. It also allows practitioners to refine sampling procedures if necessary. Monitoring during implementation helps insure that the project is being implemented as planned or if modifications need to be made.

Monitoring is an essential component of all restoration efforts. Without effective monitoring, restoration projects are exposed to several risks. For example, it may not be possible to obtain early warnings indicating that a restoration project is not on track. Without sound scientific monitoring, it is difficult to gauge how well a restoration site is functioning ecologically both

before and after implementation. Monitoring is necessary to assess whether specific project goals and objectives (both ecological and human dimensions) are being met, and to determine what measures might need to be taken to better achieve those goals. In addition, the lack of monitoring may lead to poor project coordination and decreased efficiency.

Sharing of data and protocols with others working in the same area is also encouraged. If multiple projects in the same watershed or ecosystem are not designed and evaluated using a complementary set of protocols, a disjointed effort may produce a patchwork of restoration sites with varying degrees of success (Galatowitsch et al. 1998-1999) and no way to assess system-wide progress. This would result in a decreased ability to compare results or approaches among projects.

CONTEXT AND ORGANIZATION OF INFORMATION

In 2000, Congress passed the *Estuary Restoration Act (ERA), Title I of the Estuaries and Clean Waters Act of 2000*. The ERA establishes a goal of one million acres of coastal habitats (including those of the Great Lakes) to be restored by 2010. The ERA also declares that anyone seeking funds for a restoration project needs to have a monitoring plan to show how the progress of the restoration will be tracked over time. The National Oceanic and Atmospheric Administration (NOAA) was tasked with developing monitoring guidance for coastal restoration practitioners whether they be academics, private consultants, members of state, Tribal or local government, non-governmental organizations (NGOs), or private citizens, regardless of their level of expertise.

To accomplish this task, NOAA has provided guidance to the public in two volumes. The first, *Science-Based Restoration Monitoring of Coastal Habitats, Volume One: A Framework*

for *Monitoring Plans Under the Estuaries and Clean Waters Act of 2000 (Public Law 160-457)* was released in 2003. It outlines the steps necessary to develop a monitoring plan for any coastal habitat restoration project. *Volume One* briefly describes each of the habitats covered and provides three matrices to help practitioners choose which habitat characteristics may be most appropriate to monitor for their project. Experienced restoration practitioners, biologists, and ecologists as well as those new to coastal habitat restoration and ecology can benefit from the step-by-step approach to designing a monitoring plan outlined in *Volume One*.

Volume Two, Tools for Monitoring Coastal Habitats expands upon the information in *Volume One* and is divided into two sections **Monitoring Progress Toward Goals** (Chapters 2-14) and **Context for Restoration** (Chapters 15-18). The first section, Monitoring Progress Toward Goals includes:

- Detailed information on the structural and functional characteristics of each habitat that may be of use in restoration monitoring
- Annotated bibliographies, by habitat, of restoration-related literature and technical methods manuals, and
- A chapter discussing many of the human dimensions aspects of restoration monitoring

The second section, Context for Restoration includes:

- A review of methods to select reference conditions
- A sample list of costs associated with restoration and restoration monitoring
- An overview of an online, searchable database of coastal monitoring projects from around the United States, and
- A review of federal legislation that supports restoration and restoration monitoring

The Audience

Volumes One and Two of Science-Based Restoration Monitoring of Coastal Habitats are written for those involved in developing and implementing restoration monitoring plans, both scientists and non-scientists alike. The intended audience includes restoration professionals in academia and private industry, as well as those in Federal, state, local, and Tribal governments. Volunteer groups, non-governmental organizations, environmental advocates, and individuals participating in restoration monitoring planning will also find this information valuable. Whereas *Volume One* is designed to be usable by any restoration practitioner, regardless of their level of expertise, *Volume Two* is designed more for practitioners who do not have extensive experience in coastal ecology. Seasoned veterans in coastal habitat ecology, however, may also benefit from the annotated bibliographies, literature review, and other tools provided.

The information presented in *Volume Two* is not intended as a 'how to' or methods manual: many of these are already available on a regional or habitat-specific basis. *Volume Two* does not provide detailed procedures that practitioners can directly use in the field to monitor habitat characteristics. The tremendous diversity of coastal habitats across the United States, the types and levels of impact to them, the differing scales of restoration activities, and variety of techniques used in restoration and restoration monitoring prevent the development of universal protocols. Thus, the authors have taken the approach of explaining *what one can measure during restoration monitoring, why it is important, and what information it provides* about the progress of the restoration effort. The authors of each chapter also believe that monitoring plans must be derived from the goals of the restoration project itself. Thus, each monitoring effort has the potential to be

unique. The authors suggest, however, that restoration practitioners seek out the advice of regional experts, share data, and use similar data collection techniques with others in their area to increase the knowledge and understanding of their local and regional habitats. The online database of monitoring projects described in Chapter 17 is intended to facilitate this exchange of information.

The authors do not expect that every characteristic and parameter described herein

will be measured, in fact, very few of them will be as part of any particular monitoring effort. A comprehensive discussion of all potential characteristics is, however, necessary so that practitioners may choose those that are most appropriate for their monitoring program. In addition, although the language used in *Volume Two* is geared toward restoration monitoring, the characteristics and parameters discussed could also be used in ecological monitoring and in the selection of reference conditions as well.

MONITORING PROGRESS TOWARDS GOALS

The progress of a restoration project can be monitored through the use of traditional ecological characteristics (Chapters 2 - 13) and/or emerging techniques that incorporate human dimensions (Chapter 14).

THE HABITAT CHAPTERS

Thirteen coastal habitats are discussed in twelve chapters. Each chapter follows a format that allows users to move directly to the information needed, rather than reading the whole text as one would a novel. There is, however, substantial variation in the level of detail among the chapters. The depth of information presented reflects the extent of restoration, monitoring, and general ecological literature associated with that habitat. That is, some habitats such as marshes, SAV, and oyster reefs have been the subject of extensive restoration efforts, while others such as rocky intertidal and rock bottom habitats have not. Even within habitats there can be considerable differences in the amount of information available on various structural and functional characteristics and guidance on selecting parameters to measure them. The information presented for each habitat has been derived from extensive literature reviews of restoration and ecological monitoring studies. Each habitat chapter was then reviewed by experts for content to ensure that the information provided represented the most current scientific understanding of the ecology of these systems as it relates to restoration monitoring.

Habitat characteristics are divided into two types: structural and functional. Structural habitat characteristics define the physical composition of a habitat. Examples of structural characteristics include:

- Sediment grain size
- Water source and velocity

- Depth and timing of flooding, and
- Topography and bathymetry

Structural characteristics such as these are often manipulated during restoration efforts to bring about changes in function. Functional characteristics are the ecological services a habitat provides. Examples include:

- Primary productivity
- Providing spawning, nursery, and feeding grounds
- Nutrient cycling, and
- Floodwater storage

Structural characteristics determine whether or not a particular habitat is able to exist in a given area. They will often be the first ones monitored during a restoration project. Once the proper set of structural characteristics is in place and the biological components of the habitat begin to become established, functional characteristics may be added to the monitoring program. Although structural characteristics have historically been more commonly monitored during restoration efforts, measurements of functional characteristics provide a better estimate of whether or not a restored area is truly performing the economic and ecological services desired. Therefore, incorporating measurements of functional characteristics in restoration monitoring plans is strongly encouraged.

When developing a restoration monitoring plan, practitioners should follow the twelve-step process presented in *Volume One* and refer to the appropriate chapters in *Volume Two* (habitat and human dimensions) to assist them in selecting characteristics to monitor. The information presented in the habitat chapters is derived from and expands upon the *Volume One* matrices (*Volume One* Appendix II).

Organization of Information

Each of the habitat chapters is structured as follows:

1. Introduction
 - a. Habitat description and distribution
 - b. General ecology
 - c. Human impacts to the habitat
2. Structural and functional characteristics
 - a. Each structural and functional characteristic identified for the habitat in the *Volume One* matrices is explained in detail. Structural and functional characteristics have generally been discussed in separate sections of each chapter. Occasionally, some functions are so intertwined with structural characteristics that the two are discussed together.
 - b. Whenever possible, potential methods to measure, sample, and/or monitor each characteristic are introduced or readers are directed to more thorough sources of information. In some cases, not enough information was found while reviewing the literature to make specific recommendations. In these cases, readers are encouraged to use the primary literature cited within the text for methods and additional information.
3. Matrices of the structural and functional characteristics and parameters suggested for use in restoration monitoring
 - a. These two matrices are habitat-specific distillations of the *Volume One* matrices
 - b. Habitat characteristics are cross-walked with parameters appropriate for monitoring change in that characteristic. Parameters include both those that are direct measures of a particular characteristic as well as those that are indirectly related and may influence a particular characteristic or related parameter. Tables 1 and 2 can be used to illustrate an example. The parameter of salinity in submerged aquatic

vegetation is a direct measure of a structural characteristic (salinity, Table 1). In addition, salinity is related to other structural characteristics such as tides and water source. Salinity is also related to functional characteristics such as biodiversity and nutrient cycling and may be appropriate to include in the monitoring of these functions as well (Table 2). Experienced practitioners will note that many characteristics and parameters may be related to one another but are not shown as such in a particular matrix. The matrices are not intended to be all inclusive of each and every possible interaction. The matrices provided and the linkages illustrated are only intended as starting points in the process of developing lists of parameters that may be useful in measuring particular characteristics and understanding some of their interrelationships.

- c. Some parameters and characteristics are noted as being highly recommended for any and all monitoring efforts as they represent critical components of the habitat while others may or may not be appropriate for use depending on the goals of the individual restoration project.
4. Acknowledgement of reviewers
5. Literature Cited

Three appendices are also provided for each habitat chapter. In the online form of *Volume Two*, these appendices download with the rest of the habitat chapter text. In the printed versions of *Volume Two*, each chapter's appendices are provided on a searchable CD-ROM located inside the back cover. Each appendix is organized as follows:

Appendix I - An Annotated Bibliography

- a. Overview of case studies of restoration monitoring and general ecological studies pertinent to restoration monitoring
- b. Entries are alphabetized by author

Parameters to Monitor the Structural Characteristics of SAV (excerpt)

Parameters to Monitor		Biological				Physical				Hydrological				Chemical	
Salinity (in tidal areas)		Habitat created by plants				Sediment grain size ¹				Topography / Bathymetry				Turbidity	

Appendix II - Review of Technical and Methods Manuals

These include reviews of:

- a. Restoration manuals
- b. Volunteer monitoring protocols
- c. Lab methods
- d. Identification keys, and
- e. Sampling methods manuals

Whenever possible, web addresses where these resources can be found free of charge are provided.

Appendix III - Contact information for experts who have agreed to be contacted with questions from practitioners

As extensive as these resources are, it is inevitable that some examples, articles, reports, and methods manuals have been omitted. Therefore, these chapters should not be used in isolation. Instead, they should be used as a supplement to and extension of:

- The material presented in *Volume One*
- Resources provided in the appendices
- The advice of regional habitat experts, and
- Research on the local habitat to be restored

WHAT ARE THE HABITATS?

The number and type of habitats available in any given estuary is a product of a complex mixture of the local physical and hydrological characteristics of the water body and the organisms living there. The ERA Estuary Habitat Restoration Strategy (Federal Register 2002) dictates that the Cowardin et al. (1979) classification system should be followed in organizing this restoration monitoring information. The Cowardin system is a national

standard for wetland mapping, monitoring, and data reporting, and contains 64 different categories of estuarine and tidally influenced habitats. Definitions, terminology, and the list of habitat types continue to increase in number as the system is modified. Discussion of such a large number of habitat types would be unwieldy. The habitat types presented in this document, therefore, needed to be smaller in number, broad in scope, and flexible in definition. The 13 habitats described in this document are, however, generally based on that of Cowardin et al. (1979).

Restoration practitioners should consider local conditions within their project area to select which general habitat types are present and which monitoring measures might apply. In many cases, a project area will contain more than one habitat type. To appropriately determine the habitats within a project area, the practitioner should gather surveys and aerial photographs of the project area. From this information, he or she will be able to break down the project area into a number of smaller areas that share basic structural characteristics. The practitioner should then determine the habitat type for each of these smaller areas. For example, a practitioner working in a riparian area may find a project area contains a *water column*, *riverine forest*, *rocky shoreline*, and *rock bottom*. Similarly, someone working to restore an area associated with a tidal creek or stream may find the project area contains *water column*, *marshes*, *soft shoreline*, *soft bottom*, and *oyster beds*. Virtually all estuary restoration projects will incorporate characteristics of the water column. Therefore, all practitioners should read *Chapter 2: Restoration Monitoring of the Water Column* in addition to any additional chapters necessary.

Habitat Decision Tree

A Habitat Decision Tree has been developed to assist in the easy differentiation among the habitats included in this manual. The decision tree allows readers to overcome the restraints of varying habitat related terminology in deciding which habitat definitions best describe those in their project area. Brief definitions of each habitat are provided at the end of the key.

1. a. Habitat consists of open water and does not include substrate (**Water Column**)
b. Habitat includes substrate (go to 2)
2. a. Habitat is continually submerged under most conditions (go to 3)
b. Habitat substrate is exposed to air as a regular part of its hydroperiod (go to 8)
3. a. Habitat is largely unvegetated (go to 4)
b. Habitat is dominated by vegetation (go to 7)
4. a. Substrate is composed primarily of soft materials, such as mud, silt, sand, or clay (**Soft Bottom**)
b. Substrate is composed primarily of hard materials, either of biological or geological origin (go to 5)
5. a. Substrate is composed of geologic material, such as boulders, bedrock outcrops, gravel, or cobble (**Rock Bottom**)
b. Substrate is biological in origin (go to 6)
6. a. Substrate was built primarily by oysters, such as *Crassostrea virginica* (**Oyster Reefs**)
b. Substrate was built primarily by corals (**Coral Reefs**)
7. a. Habitat is dominated by macroalgae (**Kelp and Other Macroalgae**)
b. Habitat is dominated by rooted vascular plants (**Submerged Aquatic Vegetation - SAV**)
8. a. Habitat is not predominantly vegetated (go to 9)
b. Habitat is dominated by vegetation (go to 10)
9. a. Substrate is hard, made up materials such as bedrock outcrops, boulders, and cobble (**Rocky Shoreline**)
b. Substrate is soft, made up of materials such as sand or mud (**Soft Shoreline**)
10. a. Habitat is dominated by herbaceous, emergent, vascular plants. The water table is at or near the soil surface or the area is shallowly flooded (**Marshes**)
b. Habitat is dominated by woody plants (go to 11)
11. a. The dominant woody plants present are mangroves, including the genera *Avicennia*, *Rhizophora*, and *Laguncularia* (**Mangrove Swamps**)
b. The dominant woody plants are other than mangroves (go to 12)
12. a. Forested habitat experiencing prolonged flooding, such as in areas along lakes, rivers, and in large coastal wetland complexes. Typical dominant vegetation includes bald cypress (*Taxodium distichum*), black gum (*Nyssa sylvatica*), and water tupelo (*Nyssa aquatica*). (**Deepwater Swamps**)
b. Forested habitat along streams and in floodplains that do not experience prolonged flooding (**Riverine Forests**)

Water column - A conceptual volume of water extending from the water surface down to, but not including the substrate. It is found in marine, estuarine, river, and lacustrine systems.

Rock bottom - Includes all wetlands and deepwater habitats with substrates having an aerial cover of stones, boulders, or bedrock 75% or greater and vegetative cover of less than 30% (Cowardin et al. 1979). Water regimes are restricted to subtidal, permanently flooded, intermittently exposed, and semi-permanently flooded. The rock bottom habitats addressed in *Volume Two* include bedrock and rubble.

Coral reefs - Highly diverse ecosystems, found in warm, clear, shallow waters of tropical oceans worldwide. They are composed of marine polyps that secrete a hard calcium carbonate skeleton, which serves as a base or substrate for the colony.

Oyster reefs - Dense, highly structured communities of individual oysters growing on the shells of dead oysters.

Soft bottom - Loose, unconsolidated substrate characterized by fine to coarse-grained sediment.

Kelp and other macroalgae - Relatively shallow (less than 50 m deep) subtidal and intertidal algal communities dominated by very large brown algae. Kelp and other macroalgae grow on hard or consolidated substrates forming extensive three-dimensional structures that support numerous plant and animal communities.

Rocky shoreline - Extensive littoral habitats on high-energy coasts (i.e., subject to erosion from waves) characterized by bedrock, stones, or boulders with a cover of 75% or more and less than 30% cover of vegetation. The substrate is, however, stable enough to permit the attachment and growth of sessile or sedentary invertebrates and attached algae or lichens.

Soft shoreline - Unconsolidated shore includes all habitats having three characteristics:

(1) unconsolidated substrates with less than 75% aerial cover of stones, boulders, or bedrock; (2) less than 30% aerial cover of vegetation other than pioneering plants; and (3) any of the following water regimes: irregularly exposed, regularly flooded, irregularly flooded, seasonally flooded, temporarily flooded, intermittently flooded, saturated, or artificially flooded (Cowardin et al. 1979). This definition includes cobble-gravel, sand, and mud. However, for the purpose of this document, cobble-gravel is not addressed.

Submerged aquatic vegetation (SAV; includes marine, brackish, and freshwater) - Seagrasses and other rooted aquatic plants growing on soft sediments in sheltered shallow waters of estuaries, bays, lagoons, rivers, and lakes. Freshwater species are adapted to the short- and long-term water level fluctuations typical of freshwater ecosystems.

Marshes (marine, brackish, and freshwater) - Transitional habitats between terrestrial and aquatic systems where the water table is usually at or near the surface, or the land is covered by shallow water tidally or seasonally. Freshwater species are adapted to the short- and long-term water level fluctuations typical of freshwater ecosystems.

Mangrove swamps - Swamps dominated by shrubs (*Avicenna*, *Rhizophora*, and *Laguncularia*) that live between the sea and the land in areas that are inundated by tides. Mangroves thrive along protected shores with fine-grained sediments where the mean temperature during the coldest month is greater than 20° C; this limits their northern distribution.

Deepwater swamps - Forested wetlands that develop along edges of lakes, alluvial river swamps, in slow-flowing strands, and in large coastal-wetland complexes. They can be found along the Atlantic and Gulf Coasts and throughout the Mississippi River valley.

They are distinguished from other forested habitats by the tolerance of the dominant vegetation to prolonged flooding.

Riverine forests - Forests found along sluggish streams, drainage depressions, and in large alluvial floodplains. Although associated with deepwater swamps in the southeastern United States, riverine forests are found throughout the United States in areas that do not have prolonged flooding.

THE HUMAN DIMENSIONS CHAPTER

The discussion of human dimensions helps restoration practitioners better understand how to select measurable objectives that allow for the appropriate assessment of the benefits of coastal restoration projects to human communities and economies. Traditionally, consideration of human dimensions issues has not been included as a standard component of most coastal restoration projects. Most restoration programs do not currently integrate social or economic factors into restoration monitoring, and few restoration projects have implemented full-scale human dimensions monitoring. Although some restoration plans are developed in an institutional setting that require more deliberate consideration of human dimensions impacts and goals, this does not generally extend to the monitoring stage. It is becoming increasingly evident, however, that decisions regarding restoration cannot be made solely by using ecological parameters alone but should also involve considerations of impacts on and benefits to human populations, as well. Local communities have a vested interest in coastal restoration and are directly impacted by the outcome of restoration projects in terms of aesthetics, economics, or culture. Human dimensions goals and objectives whether currently available or yet to be developed should reflect societal uses and values of the resource to be restored. Establishing these types of parameters will increase the public's understanding of the potential benefits of a

restoration project and will increase public support for restoration activities.

While ecologists work to monitor the restoration of biological, physical, and chemical functional characteristics of coastal ecosystems, human dimensions professionals identify and describe how people value, utilize, and benefit from the restoration of coastal habitats. The monitoring and observation of coastal resource stakeholders allows us to determine who cares about coastal restoration, why coastal restoration is important to them, and how coastal restoration changes people's lives. The human dimensions chapter will help restoration practitioners identify:

- 1) Human dimensions goals and objectives of a project
- 2) Measurable parameters that can be monitored to determine if those goals are being met, and
- 3) Social science research methods, techniques, and data sources available for monitoring these parameters

This chapter includes a discussion of the diverse and dynamic social values that people place on natural resources, and the role these values play in natural resource policy and management. Additionally, some of the general factors to consider in the selection and monitoring of human dimensions goals/objectives of coastal restoration are presented, followed by a discussion of some specific human dimensions goals, objectives, and measurable parameters that may be included in a coastal restoration project. An annotated bibliography of key references and a matrix of human dimensions goals and measurable parameters are provided as appendices at the end of this chapter. Also included, as an appendix, is a list of human dimensions research experts (and their areas of expertise) that you may contact for additional information or advice.

CONTEXT FOR RESTORATION

The final four chapters of this manual are designed to provide readers with additional information that should enhance their ability to develop and carry out strong restoration monitoring plans. Chapter 15 reviews methods available for choosing areas or conditions to which a restoration site may be compared both for the purpose of setting goals during project planning and for monitoring the development of the restored site over time. Chapter 16 is a listing of generalized costs of personnel, labor, and equipment to assist in the development of planning preliminary cost estimates of restoration monitoring activities. Some of this information will also be pertinent to estimating costs of implementing a restoration project as well. Chapter 17 provides a brief description of the online review of monitoring programs in the United States. The database can be accessed through the NOAA Restoration Portal (<http://restoration.noaa.gov/>). This database will allow interested parties to search by parameters and methodologies used in monitoring, find and contact responsible persons, and provide examples that could serve as models for establishment or improvement of their own monitoring efforts. Chapter 18 is a summary of the major United States Acts that support restoration monitoring. This information will provide material important in the development of a monitoring plan. A Glossary of many scientific terms is also provided at the end of the document.

References

- Cowardin, L. M., V. Carter, F. C. Golet and E. T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States, 104 pp. FWS/OBS-79/31, U.S. Fish and Wildlife Service, Washington, D.C.
- Dahl, T. E. 1990. Wetland loss in the United States 1780's to 1980's, United States Department of Interior, Fish and Wildlife Service, Washington, D.C.
- ERA. 2000. Estuary Restoration Act of 2000: Report (to accompany H.R. 1775) (including cost estimate of the Congressional Budget Office). Corp Author(s): United States. Congress. House. Committee on Transportation and Infrastructure. U.S. G.P.O., Washington, D.C.
- Federal Register. 2002. Final estuary habitat restoration strategy prepared by the estuary habitat restoration council. December 3. 71942-71949.
- Galatowitsch, S. M., D. C. Whited and J. R. Tester. 1998-1999. Development of community metrics to evaluate recovery of Minnesota wetlands. *Journal of Aquatic Ecosystem Stress and Recovery* 6:217-234.
- Keddy, P. A. 2000. Wetland Ecology: Principles and Conservation. Cambridge University Press, Cambridge, United Kingdom.
- Meeker, S., A. Reid, J. Schloss and A. Hayden. 1996. Great Bay Watch: A Citizen Water Monitoring Programpp. UNMP-AR-SG96-7, University of New Hampshire/University of Maine Sea Grant College Program.
- Mitsch, W. J. and J. G. Gosselink. 2000. Wetlands. Third ed. Van Nostrand Reinhold, New York, NY.
- NOAA, Environmental Protection Agency, Army Corps of Engineers, United States Fish and Wildlife Service and Natural Resources Conservation Service. 2002. An Introduction and User's Guide to Wetland Restoration, Creation, and Enhancement (pre-print copy), Silver Spring, MD.
- Washington, H., J. Malloy, R. Lonie, D. Love, J. Dumbrell, P. Bennett and S. Baldwin. 2000. Aspects of Catchment Health: A Community Environmental Assessment and Monitoring Manual. Hawkesbury-Nepean Catchment Management Trust, Windsor, Australia.